

Over the age of 65 with A-Fib and not on an Anticoagulant? You may be eligible to join a Pioneering Clinical Trial!



Testing Abrelacimab, an Investigational, Monthly Injectable for
Stroke Prevention in A-Fib Patients



Goals of the Study

The **LILAC - TIMI 76** trial is looking for participants who have non-valvular atrial fibrillation (AF), including paroxysmal AF, and have been told by their doctors that they can't take oral blood-thinning medication because they have a high risk of bleeding.

Who May Participate?



History of atrial fibrillation

Current episode not necessary



Not on a current blood thinner like Apixaban, Rivaroxaban, or Warfarin in the past 60 days

Be able to explain why you are unable to be on an oral blood thinner



If left atrial appendage closure (**Watchman Device**) isn't right for you, consider the LILAC - TIMI 76 trial



AFIB and Stroke

#1 preventable cause of stroke is AF



5x stroke risk



↑ 2x mortality

~90%

 patients require life-long anticoagulation

If interested, please call or email a member of the LILAC study team:

[Placeholder for contact info]

*67% of patients with newly diagnosed AF in 2020 were not started on an anticoagulant within 12 months of diagnosis (Ko et al. (2022) JAMA). Reasons you may not be prescribed an anticoagulant include: Older age, fall risk, dementia, frailty, anaemia, prior hip/pelvic fracture, chronic kidney disease, or personal choice (LILAC_Poster_English UK_V2_12Oct2023)